

# Nutrition for distance events

Louise M Burke<sup>1</sup>, Gregoire Millet<sup>2</sup>, Mark A Tarnopolsky<sup>3</sup>

<sup>1</sup>Department of Sports Nutrition, Australian Institute of Sport, Belconnen, Australia 2616

<sup>2</sup> Department of performance enhancement, Aspire, PO Box 22287, Doha, Qatar

<sup>3</sup> Departments of Pediatrics and Medicine, McMaster University, Hamilton, Canada.

Corresponding author:

Louise M Burke,  
Department of Sports Nutrition,  
Australian Institute of Sport,  
Belconnen,  
Australia 2616

Ph 61 2 6214 1351

Fax 61 2 6214 1603

Email: [louise.burke@ausport.gov.au](mailto:louise.burke@ausport.gov.au)

Running title: Nutrition for distance events

## **Abstract:**

The goal of training is to prepare the distance athlete to perform at his or her best during major competitions. Whatever the event, nutrition plays a substantial role in the achievement of various factors that will see a runner or walker take the starting line in the best possible form. Everyday eating patterns must supply fuel and nutrients needed to optimise their performance during training sessions and to recover quickly afterward. Carbohydrate and fluid intake before, during, and after a workout may help to reduce fatigue and enhance performance. Recovery eating should also consider issues for adaptation and the immune system which may involve intakes of protein and some micronutrients. Race preparation strategies should include preparation of adequate fuel stores, including carbohydrate loading for prolonged events such as the marathon or 50 km walk. Fluid and carbohydrate intake during races lasting an hour or longer should also be considered. Sports foods and supplements of value to distance athletes include sports drinks and liquid meal supplements to allow nutrition goals to be achieved when normal foods are not practical. While caffeine is an ergogenic aid of possible value to distance athletes, most other supplements are of minimal benefit.

## Introduction

There are a large number of events which involve prolonged effort within the IAAF umbrella of track and field, road running, cross country, and race-walking. Events commonly undertaken by elite competitors include the 5,000 and 10,000 m track events, the half-marathon and marathon, the 20 and 50 km walks and cross-country runs (8 km for females and 12 km for males). In addition, there is a vast array of “fun runs” and community events around the world which can attract large fields ranging from the elite to the weekend warrior. Nutrition plays a key role in assisting distance athletes of all levels to achieve their training and competition goals.

## Training for distance runners and walkers

Distance runners follow a periodised training program (see Stellingwerff et al., 2007), split into base training (8 to 16 weeks), a precompetitive period (8-6 weeks) and a competitive period (if track events) or a tapering phase (up to 3 weeks) prior to a marathon followed by a short transition/recovery phase. Heat acclimatisation before competition in a hot environment and altitude training are other specialised training techniques often undertaken by distance runners and walkers. Altitude training remains a controversial area, with coaches and scientists still arguing over the benefits of periods in a hypoxic (lower oxygen) environment on performance at sea level. Distance athletes who usually reside at low altitudes have a variety of options for undertaking altitude training (see Hawley et al., 2007).

Since the athletes of East Africa origins (Kenya, Ethiopia and more recently, Eritrea and Uganda) dominate distance running, the reasons for their superiority have been extensively studied (Billat et al., 2003; Lucia et al., 2007; Saltin et al. 1995). Among the key physiological factors of distance running performance (i.e.  $VO_{2max}$ , the maximal fraction of  $VO_{2max}$  sustained during the event, the velocity at lactate threshold and the running energy cost), it appears that they have mainly a greater running economy (Lucia et al., 2006) and a higher fractional utilisation of  $VO_{2max}$  than Caucasian runners (Saltin et al., 1995; Lucia et al., 2006). The underlying mechanisms of these differences are still contradictory but are a combination of social, genetic and anthropometric/biomechanical factors. The effects of the altitude residence/training *per se* or of nutritional differences have not been identified. Although a high running distance in training (80-150 km.wk<sup>-1</sup> in 3,000–5,000 m runners; 150-220 km.wk<sup>-1</sup> in marathon runners during base training) is commonly observed in all distance runners, several studies have reported that most African runners spend a larger part of their weekly training at high relative intensity. The current trend in distance running is to a ‘*polarised training*’ model; i.e. a large percentage (70-75%) at strictly aerobic intensity, a small percentage (<10%) of ‘Tempo’ training at around or above lactate threshold and 15-20% at high intensity.

## Competition in distance events

Although most distance events involve a single race, there are some events which require heats and finals (e.g. 5,000 and 10,000 m on the Olympic Games and World Championship programs). Most distance runners and walkers peak for several important events in a year (e.g. a Big City Marathon or the World Championships). However, there may be other situations in which they compete in a series of races including the lucrative professional circuit in Europe, meets within a university circuit such as the NCAA season, or in the cross-country schedule for club athletes. In general, the main competition for track and field occurs in summer, whereas cross-country has an autumn and/or winter season. Most road races attracting large fields of both elite and community-based participants are scheduled over the warmer months from spring to late autumn where heat and hydration become more of an issue. The schedule of Big City Marathons, which includes races in

Boston, Chicago, New York, London, and Paris, extends from April to November.

Aerobic metabolism typically accounts for more than 95% of the energy cost of long-distance events, especially half-marathon and marathon races and the longer walking events. However, there are critical times in all distance races requiring anaerobic effort—for example, a surge, a hill, or a sprint finish—that may be the ultimate factor in determining the order of race finishers. The factors that limit the performance of distance runners and walkers vary according to the duration and environment of the race and nutrition is an important factor in success over the duration of the event. Because many of these factors—such as fluid balance, the availability of carbohydrate fuel, and even the disturbance to acid-base status arising from anaerobic glycolysis—can be manipulated by dietary strategies, nutrition is an important component of the athlete's preparation for competition.

### **Nutritional Issues and Challenges**

There is a range of common nutritional issues that arise in long-distance running and walking as a result of issues related to optimal physique, training, and race day performance. This review will provide an overview of the major issues.

#### ***Physique***

Very low levels of body fat are a striking feature of successful distance athletes. However, it is hard to distinguish whether this is a critical factor in determining successful performance or the outcome of the high training volumes needed for successful performance. Low levels of both total mass (which determines the total energy cost of running) and fat mass (dead weight that must be transported) assist fast and economical movement. These traits become even more important when the event involves long distances or moving against gravity (e.g., running up hills in a road or cross country race). Because upper-body musculature is unimportant for running performance, elite runners and walkers typically exhibit minimal evidence of muscle development in their arms and upper torso. Although there is variability in the size of long-distance runners and walkers, the winners of "hot weather" races tend to be small and light. A small and compact physique offers thermoregulatory advantages, both by reducing the absolute amount of heat that is produced (smaller muscle mass) and by achieving a more efficient dissipation of heat generated by the body (enhanced ratio of surface area to volume). There are data from both modelling (Dennis and Noakes, 1999) and laboratory (Marino et al., 2000) sources to show that lighter runners store less heat at the same running speed and enjoy an advantage in conditions where heat dissipation mechanisms are at their limit.

Some runners and walkers achieve a small and very lean frame as a result of their genetic background and training program. However, other runners with naturally larger frames or higher levels of adiposity feel that they must whittle themselves down to an "unnatural" size and body fat level to be competitive. Although many male runners eat and train specifically to reduce their body fat and racing weight, the battle for a low percentage of body fat and weight control is most often identified as a female problem. This may be because females generally need to push their body characteristics further from their natural shape than male runners to achieve the leanness that is considered ideal. Attempts to deviate body fat further from the apparent biological "default" can have negative effects, including "penalties" resulting from the low body fat levels *per se* such as a lack of insulation against cold. Other penalties arise from the nutrition and training methods used to manipulate weight and body fat including restricted intakes of energy, protein, carbohydrate and micronutrients (Burke, 2007). Some athletes develop frank medical or psychiatric problems such as eating disorders, osteopaenia and

chronic menstrual dysfunction. More develop sub-clinical versions of these problems; the spectrum of restrained eating, menstrual dysfunction and poor bone health within the "female athlete triad" is covered in greater detail by Manore et al. (2007) and similar issues should also be considered in the evaluation of some male athletes.

The problems associated with poor bone health lie not only with the risk of a premature onset of osteoporosis but also with the immediate problem of stress fractures. Recurrent or chronic stress fractures can prevent the athlete from competing at important times and interfere with his or her ability to undertake the training volume necessary for high level performance. Many athletes have had promising careers ended by this injury pattern. Distance runners and walkers should be encouraged to set realistic weight and body fat goals; these are specific to each athlete and must be judged by trial and error over a period of time. Further discussion on dietary strategies to assist with loss of weight and body fat is found in the review by O'Connor et al. (2007).

### ***Poor iron status***

There is a common belief that endurance athletes, particularly distance runners, are at high risk of iron deficiency; this has been given apparent credibility because the target levels for iron status measures such as serum ferritin are often set well above those of normal population standards to provide a "safety margin" for athletes whose performance are underpinned by the roles of iron in oxygen transport (haemoglobin and myoglobin) and enzyme function (for review, see Deakin 2006).

The depletion of the body's iron stores progresses through a number of stages with different functional and diagnostic criteria (see Deakin 2006). The literature is unclear, in part because of methodological concerns, whether iron depletion, in the absence of anaemia impairs exercise performance (Fogelholm 1995). Some studies of iron supplementation in iron-depleted but non-anaemic female runners (Klingshirn et al. 1992; Newhouse et al. 1989; Powell and Tucker 1991) failed to find differences in performance changes between supplementation and placebo treatment groups, even when serum ferritin increased with iron therapy (Klingshirn et al. 1992; Newhouse et al. 1989). However, in other studies, female runners with low ferritin levels experienced a performance improvement, albeit in conjunction with an increase in haemoglobin levels, after iron supplementation (Lamanca and Haymes 1993; Schoene et al. 1983). Of course, athletes are also concerned whether iron depletion affects their ability to recover between workouts or races. Brownlie et al. (2004) exposed previously untrained participants with non-anaemic iron depletion to a 4-week training program and found that those with a tissue iron deficiency (based on abnormal serum transferrin receptor concentrations) had an impaired adaptation to this training compared with a similar group who received iron supplements. In contrast, iron supplementation did not affect endurance cycling performance at the end of the training program in the iron-depleted group who were not tissue iron-depleted.

In summary, the true prevalence of iron-deficiency anaemia in distance runners and walker is probably not greater than in the general population (Fogelholm, 1995). However, reduced iron status does occur and may be problematic for performance or adaptation to training, particularly altitude training (see Hawley et al., 2007). The cause is essentially the same reason that it occurs in the general population: a lower than desirable intake of high bioavailability iron. Iron requirements may be increased in distance athletes because of increased gastrointestinal or haemolytic iron losses (for review see Deakin, 2006). However, the most important risk factor is still the low-energy or low-iron diet. Females, vegetarian eaters, and those following diets

with restricted quantity and variety are at highest risk. Dietary interventions to reverse or prevent a decline in iron status involve strategies to increase total iron intake as well as to increase the bioavailability of this iron.

The management and prevention of iron deficiency requires careful diagnosis using a variety of clinical, haematological, dietary, and medical data. Haematological and biochemical tests that are routinely measured to indicate iron status should be undertaken in a way that minimises or standardises the effect of exercise on the results. In athletic populations, ferritin levels lower than 30-35 ng/ml (Nielsen and Nachtigall 1998) are generally marked for further consideration or review, especially where it makes a change in the established iron status history of the individual. New tests including the measurement of serum transferrin receptors and the characteristics of reticulocytes may offer new opportunities. However, these tests are not routinely available in all laboratories and need to be studied carefully in relation to iron status in athletes.

Many distance athletes are tempted to self-medicate with iron supplements that can be purchased over the counter. However, there are several risks involved with the consumption of iron supplements in the absence of a confirmed iron status problem, including haemosiderosis or iron overload. Typically, a 3-month period of supplementation, in the form of a daily dose of 100 mg of elemental iron, is needed to restore depleted iron stores (Nielsen and Nachtigall 1998). In some cases, when it is not possible to enhance dietary iron intake sufficiently, iron supplementation is continued at a lower dose to prevent ongoing iron drain. In cases of extreme iron depletion or where oral iron intake is not tolerated, intramuscular injections of iron can achieve a rapid increase in iron stores. However, there is no evidence of additional performance benefits over oral supplementation, and there are higher risks of side effects. Iron injections will not increase haemoglobin levels or other iron parameters in people who are not otherwise suboptimal in iron status (Ashenden et al. 1998).

### ***Carbohydrate needs for optimal training and recovery***

Distance runners and walkers must be able to rapidly recover their muscle fuel stores between daily or twice-daily sessions, and between races on the competition circuit. A high-carbohydrate intake enhances the performance of a single bout of prolonged running as well as the recovery and performance of a subsequent running bout (Fallowfield and Williams, 1993). However, muscle glycogen concentrations may not completely recover over 24 to 48 h following a very strenuous running session (e.g., marathon) or unaccustomed eccentric loading, despite a plentiful carbohydrate supply (Asp et al. 1997; Sherman et al. 1983). Unaccustomed muscle damage may cause a disruption to muscle cell function and may require an increase in total carbohydrate intake in the first 24 h of recovery (Doyle et al. 1993) or a greater recovery time (up to 7 days) for full replacement of muscle glycogen.

Logically, benefits from enhancing acute recovery between sessions should translate over time into better training adaptations and long-term performance gains. However, the literature, which includes three studies involving runners, is curiously unclear in showing that high carbohydrate diets provide superior training outcomes to moderate carbohydrate intakes (Burke, 2007). Kirwan and colleagues (1988) studied well-trained runners who increased their training by 150% for 5 days while consuming either high (8 g/kg/d) or moderate (4 g/kg/d) intakes of carbohydrate. Muscle glycogen concentration gradually declined in both treatments but was better preserved with the higher carbohydrate diet; additionally, running economy at two different running speeds was better. By contrast Sherman and colleagues (1993) followed 7 days of training in two groups of runners who consumed daily carbohydrate intakes of either 5 g/kg (gradually reduced muscle

glycogen levels) or 10 g/kg (maintained muscle glycogen concentrations). At the end of this period, the groups did not differ in their capacity to undertake two treadmill runs to exhaustion at 80%  $\dot{V}O_2$ max with a short recovery interval at the end of a training session.

Finally, well-trained runners undertook 7 days of intensified training supported by both moderate- (5.4 g/kg/d) and high- (8.5 g/kg/d) carbohydrate diets (Achten et al. 2004). Muscle glycogen utilisation decreased during submaximal running on the moderate-carbohydrate diet and there was a decline in speed over 8 km (treadmill) and 16 km (outdoor) time-trials. However, the high-carbohydrate treatment was associated with a smaller decrease in 8 km speed and maintenance of 16 km performances. The authors concluded that a high-carbohydrate diet reduced symptoms of overreaching in runners during intensified training compared with a moderate-carbohydrate diet but could not prevent it entirely.

An emerging interest is that of dietary periodisation – the so-called “train low, compete high” approach - in which distance athletes deliberately train with low glycogen or carbohydrate availability to enhance metabolic adaptations to the training stimulus, then replete carbohydrate to enhance their competition performance (see Hawley et al. 2007). Currently, there is inadequate scientific support to recommend that distance athletes should practice carbohydrate restriction for prolonged periods. Indeed, the potential disadvantages of this practice include an increased risk of illness and injury (see Nimmo et al., 2007) and reduced well-being or capacity to train (see Burke and Kiens, 2006). In fact, the available study supporting a “train low” approach (Hansen et al., 2005) achieved glycogen depletion for some, but not all, training sessions by manipulating the training time table rather than dietary intake. Indeed, it is likely that elite athletes spontaneously periodise carbohydrate availability within their *microcycles* of training because the practicalities of their lifestyle and training mean that some sessions are taken after an overnight fast, or without complete refuelling between workouts.

Unless more sophisticated research can identify benefits from deliberately “training low”, distance athletes should eat to promote carbohydrate availability, at least for the most important training sessions of the week. Recent recommendations for daily carbohydrate intake (Burke et al., 2004) acknowledge that fuel requirements for distance athletes differ according to body size and training loads. The targets of 7-10 g/kg/d for high volume training and 5-7 g/kg/d for more moderate exercise loads provide a general target that must be fine-tuned according to overall nutritional goals and performance feedback from each athlete. Such recommendations may be unfeasible for runners, particularly females, whose focus on low body mass and body fat levels requires energy restriction and, by association, a lower carbohydrate intake. The compromise is to periodise nutrition goals and dietary carbohydrate intakes over the season, so that lower intakes and physique goals are the priority of training periods, whereas greater carbohydrate intakes are allowed during competition preparation and recovery to maximise glycogen stores.

Although total intake of carbohydrate is probably the most important determinant of post-exercise refuelling, during periods of high volume training, the distance athlete should use other dietary strategies to promote recovery. Speedy intake of carbohydrate after exercise will maximise the period of effective refuelling time (Burke et al., 2004). Carbohydrate-rich foods in recovery snacks and meals should be chosen according to the need to meet practical challenges (e.g. finding portable foods when the athlete is “on the go”) or to meet additional nutritional goals (e.g. to provide a source of iron, protein or other nutrient need). It is probably useful to co-ingest protein with carbohydrate-rich recovery snacks. Although the effect of protein on

glycogen resynthesis is likely to be minimal in most circumstances (see Tipton et al. 2007), various issues of recovery and adaptation require protein synthesis. Indeed, in addition to refuelling, the distance athlete needs to consider a range of recovery eating goals after training and races, including rehydration (Shirreffs et al., 2007); repair and adaptation (Hawley et al., 2007) and preserving the immune system (Nimmo et al., 2007)

### ***Protein requirements during training***

Data from studies of essentially recreational exercisers have led to the belief that protein requirements are not altered by any form of physical activity. However, the high volumes of training and the training intensities possible only in elite athletes result in estimated protein requirements that are nearly twice that of sedentary individuals; i.e. 1.6 – 1.7 g/kg/d (Tarnopolsky et al. 1988; Friedman & Lemon, 1989). Even for modestly trained individuals, there is an increase in protein requirements estimated from nitrogen balance experiments (Meredith et al., 1989; Phillips et al., 1993). Although no study has specifically calculated protein requirements for elite female athletes, nitrogen balance data imply that requirements for women are about 25 % less than for men; i.e. 1.2 – 1.3 g/kg/d (Phillips et al. 1993; McKenzie et al., 2000) Most athletes will achieve these protein intakes from an everyday diet providing 10-15% of energy as protein and adequate energy. Nevertheless, it is important to evaluate protein intake on a g/kg basis as opposed to a percentage of the diet to avoid low intakes that can be seen in energy restricting athletes. A low energy intake will also negatively affect protein requirements (Calloway, 1975).

There are benefits to the timing of nutrient delivery, especially when undertaking high volumes of training. When female athletes consumed a nutritional supplement immediately after each workout during a training camp, they achieved an improvement in nitrogen balance, lesser weight loss, and improved performance on a trial completed at the end of the week than when the supplement was consumed after breakfast (Roy et al., 2002).

### ***Fuelling Up for Competition***

Preparation for racing should ensure that muscle carbohydrate stores are matched to the anticipated fuel needs of the event. For races of up to 60-90 minute duration, normalised muscle glycogen stores are adequate and can generally be achieved by 24-36 h of high carbohydrate intake. Carbohydrate loading in preparation for prolonged exercise resulted from pioneering studies undertaken in the 1960s using percutaneous biopsy techniques to examine fuel utilisation and enzyme activities in the muscle. These studies on healthy but untrained men produced the classic 7-day model to supercompensate muscle glycogen stores; a 3-4 day depletion phase of hard training and low carbohydrate intake followed by a 3-4-day loading phase of high carbohydrate intake and exercise taper (Bergstrom et al., 1967). Early field studies of prolonged running events showed that this strategy enhanced sport performance, not by allowing the athlete to run faster but by prolonging the time that race pace could be maintained (Karlsson and Saltin, 1971).

A modified version of carbohydrate-loading was developed when well-trained runners were shown to super-compensate their glycogen stores without a severe depletion or glycogen stripping phase (Sherman et al., 1981). The modified protocol, consisting simply of 3 d of high carbohydrate intake and taper, was offered as a more practical competition preparation which avoided the fatigue and complexity of the extreme diet and training requirements of the previous depletion phase. More recently, muscle glycogen concentrations were measured after 1 and 3 days of rest and a high carbohydrate intake (10 g/kg body mass per day) in well-trained male athletes (Bussau et al., 2002): this study found that optimal refuelling is

probably achieved within 36 to 48 h following the last exercise session, at least when the athlete rests and consumes adequate carbohydrate.

Theoretically, carbohydrate loading can enhance performance in distance races that would otherwise be limited by the fatigue caused by glycogen depletion. Studies in well-trained runners have failed to detect benefits of carbohydrate loading for 10 km treadmill running (Pitsiladis et al., 1996), a 20.9 km race on an indoor track (Sherman et al., 1981), and a 25 km treadmill run (Sullo et al., 1998). By contrast, carbohydrate loading has been shown to enhance performance of a 30 km cross country run (Karlsson and Saltin, 1971), a 30 km treadmill run in trained men (Williams et al., 1992), and a 25 km treadmill run in moderately trained men (Sullo et al., 1998). Typically, carbohydrate loading is associated not with an increase in overall running speed but with maintenance of race pace during the last part of the run compared with the control trial or group. Therefore, runners and walkers should consider carbohydrate loading for races of 30 km and longer.

#### ***Fat adaptation – a twist on depletion prior to carbohydrate loading***

Distance runners and walkers should have a high capacity for fat oxidation during exercise as a legacy of their training. However, this capacity can be further up-regulated by as little as 5 d of training while following a low-carbohydrate (<2.5 g/kg/d), high-fat (~65-70% of energy) diet. In trained individuals, “fat adaptation” achieves a markedly increased fat oxidation and reduced utilisation of muscle glycogen (“glycogen sparing”) during subsequent sub-maximal exercise (Burke et al., 2000). This effect persists even when followed by acute strategies to carbohydrate load, and eat carbohydrate before and during the bout (for review see Burke and Kiens 2006). Such a combination of dietary strategies would seem the perfect preparation for a marathon or distance walking event, simultaneously optimising carbohydrate stores while maximising the capacity for fat oxidation. Curiously, the effect on endurance and ultra-endurance performance is unclear (Burke and Kiens, 2006).

There is now evidence that what was initially viewed as glycogen sparing may be, in fact, a down-regulation of carbohydrate metabolism or “glycogen impairment”. Fat adaptation/ carbohydrate restoration strategies are associated with a reduction in the activity of a key enzyme regulating carbohydrate metabolism, pyruvate dehydrogenase (Stellingwerff et al., 2006). Such a change would impair rates of glycogenolysis at a time when muscle carbohydrate requirements are high. This explains the observation that when fat adaptation/carbohydrate restoration are applied to exercise protocols which mimic a real-life race – self-pacing, and the interspersing of high-intensity and moderate-intensity exercise - there is a compromised ability to performance high-intensity sprints (Havemann et al. 2006) In many endurance events, the critical activities in a race – the breakaway, the surge up a hill, or the sprint to the finish line – are all dependent on the runner’s ability to work at high intensities. With growing evidence that this critical ability may be impaired, it now seems clear that fat adaptation or pre-loading depletion strategies should not be undertaken by distance athletes.

#### ***Fluid and Fuel Intake during races***

In distance running and walking events, especially road races, a network of aid stations allows competitors to consume fluids during the race. In large community participation events, a supply of water, sport drinks, and sponges is provided, although elite competitors are usually provided with opportunities to supply their own race beverages at specially marked tables. There is still debate on the ideal hydration plans for distance events, with the observation that most top runners are conservative with fluid intake while some of the “back of the pack” participants in large community events risk serious

problems from over-consumption of fluids (Noakes, 2002; Almond et al., 2005). These issues are covered in greater detail by Shirreffs et al. (2007).

The use of carbohydrate–electrolyte drinks (sport drinks) during races of 60 min or longer provides the runner or walker with the potential to replace fluid and carbohydrate simultaneously, with the option of altering the carbohydrate concentration of the drink (typically 4-8 g/100 ml), according to the priority of rehydration or refuelling in a particular event. Sports gels and confectionery are other easily consumed sources of carbohydrate often consumed by distance athletes. There is good evidence of the benefits of carbohydrate intake during prolonged (>90 min) exercise (Hargreaves 1999), with reports dating back to the Boston marathon in the 1920s that the consumption of sweets during the race prevented hypoglycaemia and enhanced running performance (Gordon et al. 1925; Levine et al. 1924). Recent studies in which carbohydrate ingestion enhanced a running protocol include a 40 km outdoor run in the heat (Millard-Stafford et al. 1992), a 30 km road run (Tsintzas et al. 1993), a marathon run on a treadmill (Tsintzas et al. 1995) and a ~2 h treadmill protocol to exhaustion at 70% VO<sub>2</sub>max (Tsintzas et al. 1996b). The generally accepted mechanisms of performance enhancement include prevention of hypoglycaemia, sparing of liver glycogen, and provision of an additional muscle fuel substrate (Hargreaves 1999). However, in the case of running, there is some evidence of muscle glycogen sparing, at least in selected fibres (Tsintzas et al. 1996a; Tsintzas et al. 1993).

The effect of carbohydrate intake during shorter distance events is unclear with the potential mechanism of any performance enhancements being attributable to effects on the central nervous system rather than provision of muscle fuel (see Burke, 2007). One study involving a 15 km treadmill run in a hot environment found an improvement in speed over the last, self-paced portion of the run when carbohydrate was ingested immediately before and during the run compared with a placebo trial (Millard-Stafford et al. 1997). By contrast, carbohydrate intake during an 18 km run failed to enhance performance of a large group of runners or the fastest runners in the group compared with water (van Nieuwenhoven et al. 2005), and highly trained runners experienced a trivial effect on performance when carbohydrate was consumed during a half-marathon (Burke et al., 2005). Further studies are needed to determine the full range of events that might benefit from carbohydrate intake immediately before and during the race.

### ***Sex differences in nutrition strategies***

It has been assumed that dietary advice for female distance athletes would be a simple extrapolation from male athletes, scaled to their smaller size. However, numerous studies have found that females oxidise more fat and less carbohydrate than men during endurance exercise (see Tarnopolsky, 2000). An early study found that increasing dietary carbohydrate intake from 55% to 75% of habitual energy intake for 4 days neither increased glycogen storage nor enhanced cycling performance in female athletes, in stark contrast to the results seen in males (Tarnopolsky et al., 1995). Of course, the relatively low energy intake of the females limited carbohydrate intake to < 6.5 g/kg/d even in the “loading” phase. A follow-up study provided an additional trial in which 75% of a higher energy intake achieved carbohydrate intakes > 8 g/kg/d (Tarnopolsky et al., 2001). With higher carbohydrate and extra energy, females increased muscle glycogen, albeit to levels which were about 50% of the increase seen in male subjects. From a practical perspective, carbohydrate loading is of use to female athletes only if they are prepared to consume adequate energy and carbohydrate.

In contrast to the limited ability for women to carbohydrate load, the dietary recommendations for men and women with respect to sport drink consumption during exercise (Riddell et al., 2003; Wallis et al., 2006), and for

post-exercise glycogen re-synthesis (Tarnopolsky et al., 1997), appear to be similar.

### **Sport Foods and Supplements**

Many distance athletes, even at a recreational level, are consumers of sport foods and supplements. Products such as sports drinks and liquid meal supplements are specially designed to help a runner or walker meet specific needs for energy, fluid and nutrients in situations where everyday foods are not practical to eat, although the expense must be considered (see Burke, 2007). Nutritional ergogenic aids have generally been poorly tested or have failed to live up to their claims when rigorous testing has been undertaken on distance running/walking performance. The exception is caffeine, which may enhance the performance of some runners (for review, see Graham, 2001). Recent research has focused on the use of small doses of caffeine before and during endurance exercise, since the benefits appear to be similar to that achieved by larger doses of 6-9 mg/kg (see Maughan et al., 2007). Caffeine intakes of as little as 3 mg/kg have been shown to enhance running performance, including a worthwhile improvement of ~1% in an 8 km track protocol (Bridge and Jones, 2006). However, runners who were provided with very small amounts of caffeine (~ 1.3 mg/kg) during an 18 km road race did not show a detectable improvement in performance (Van Nieuwenhoven et al., 2005).

While bicarbonate supplementation is typically considered a strategy for middle distance running (see Stellingwerff et al., 2007), it has been shown to improve performance of the longer track events (e.g. 5,000 m races) (Oopik et al., 2003; 2004). Creatine loading has become synonymous with the enhancement of repeated sprint training or exercise bouts (see Tipton et al., 2007) and is typically considered inappropriate for use by distance athletes. In fact, runners recorded a slower time to complete a 6 km cross-country run after creatine supplementation (Balsom et al, 1993), presumably due to the accompanying increase in body mass. In spite of recent evidence that prior creatine loading enhances the muscle's capacity for glycogen loading or re-synthesis (Nelson et al., 2001; van Loon et al., 2004), it is likely that the increase in body mass would hinder performance in distance running events, particularly if the course is hilly. Further research is needed to test the hypothesis that glycerol hyperhydration can enhance thermoregulatory function in conditions in which thermal stress limits running performance.

Finally, the claims made in support of the majority of other supplements and compounds marketed as ergogenic aids are not supported by scientific research (see Burke, 2007). Of course, more research is needed, using rigorous control and carefully chosen protocols to test the claims for most products. In many cases, particular (proposed) ergogenic compounds that are used by distance athletes have not been appropriately tested and no further comments can be made about these products. The reader is therefore referred to the general conclusions provided by Maughan et al., 2007.

### **Summary of nutrition guidelines for distance athletes**

Clear consensus for	<ul style="list-style-type: none"><li>• Distance athletes should follow established guidelines to meet the carbohydrate needs for their training loads and to enhance recovery after each training session. These strategies are particularly important to promote performance and recovery for key training sessions.</li><li>• Distance athletes should consume sufficient carbohydrate to prepare fuel stores that are adequate for their event. Carbohydrate loading or glycogen supercompensation will be of benefit to</li></ul>
---------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	<p>longer events such as the marathon or 50 km walk. A prolonged depletion phase is unnecessary and may even impair performance.</p> <ul style="list-style-type: none"> <li>• Carbohydrate and fluid intake during an event is possible and of probable value for races lasting longer than 60 minutes. Each athlete should experiment to find a plan that is practical and provides benefits for their performance.</li> <li>• Iron deficiency may be a problem for some distance runners, but this is a diagnosis of exclusion and other causes need to be ruled out. Nutritional counseling to increase intake of bioavailable iron is an important goal of prevention and therapy.</li> <li>• Some sports supplements such as sports drinks and liquid meals may be useful in providing a practical way for distance athletes to meet their nutrition goals. Moderate doses of caffeine can provide an ergogenic benefit to distance running and may be useful for some runners.</li> </ul>
Clear consensus against	<ul style="list-style-type: none"> <li>• Distance athletes should not practise extreme levels of energy restriction to achieve loss of body weight/body fat without considering the effect on their ability to meet goals for carbohydrate, protein, iron or other nutrients. Hormonal balance, bone health and the immune system are also critically impaired by inadequate energy intakes.</li> <li>• Routine supplementation with iron or iron injections in the belief that it enhances performance should be strongly discouraged in the absence of documented iron depletion or anaemia. Supplementation in the absence of deficiency can lead to serious medical conditions such as haemosiderosis.</li> <li>• The majority of supplements that are promoted to distance athletes are unlikely to provide substantial benefits, and should not replace sound eating and training practices.</li> </ul>
Issues that are equivocal	<ul style="list-style-type: none"> <li>• It is unclear whether distance athletes will enhance adaptations and performance outcomes by undertaking deliberate strategies to restrict carbohydrate availability during training. In the real world, elite athletes will probably achieve some level of periodisation of carbohydrate status within the microcycles of their training program. Any benefits of more prolonged carbohydrate depletion need to be balanced by the possible disadvantages.</li> </ul>

## References

- Achten, J., Halson, S. H., Moseley, L., Rayson, M. P., Casey, A. and Jeukendrup, A. E. (2004). Higher dietary carbohydrate content during intensified running training results in better maintenance of performance and mood state. *Journal of Applied Physiology*, 96, 1331-1340.
- Almond, C. S. D., Shin, A. Y., Fortescue, E. B., Mannix, R. C., Wypij, D., Binstadt, B. A., Duncan, C. N., Olson, D. P., Salerno, A. E., Newburger, J. W. and Greenes, D. S. (2005). Hyponatremia among runners in the Boston marathon. *New England Journal of Medicine*, 352, 1550-1556.
- Ashenden, M. J., Fricker, P. A., Ryan, R. K., Morrison, N. K., Dobson, G. P. and Hahn, A. G. (1998). The haematological response to an iron injection amongst female athletes. *International Journal of Sports Medicine*, 19, 474-478.
- Asp, S., Rohde, T. and Richter, E. A. (1997). Impaired muscle glycogen resynthesis after a marathon is not caused by decreased muscle GLUT-4 content. *Journal of Applied Physiology*, 83, 1482-1485.
- Balsom, P. D., Harridge, S. D. R., Soderlund, K., Sjodin, B. and Ekblom, B. (1993). Creatine supplementation per se does not enhance endurance exercise performance. *Acta Physiologica Scandinavica*, 149, 521-523.
- Bergstrom, J., L. Hermansen, E. Hultman and B. Saltin. 1967. Diet, muscle glycogen and physical performance. *Acta Physiologica Scandinavica* 71: 140-150.
- Billat, V., Lepretre, P. M., Heugas, A. M., Laurence, M. H., Salim, D., and Koralsztejn, J. P. (2003). Training and bioenergetic characteristics in elite male and female Kenyan runners. *Medicine and Science in Sports and Exercise*, 35, 297-304; discussion 305-296.
- Blom, C. S. B., Costill, D. L. and Vollestad, N. K. (1987). Exhaustive running: inappropriate as a stimulus of muscle glycogen supercompensation. *Medicine and Science in Sports and Exercise*, 19, 398-403.
- Bridge, C. A. and Jones, M. A. (2006). The effect of caffeine ingestion on 8 km run performance in a field setting. *Journal of Sports Sciences*, 24, 433-439.
- Brownlie, T., Utermohlen, V., Hinton, P. S. and Haas, J. D. (2004). Tissue iron deficiency without anemia impairs adaptations in endurance capacity after aerobic training in previously untrained women. *American Journal of Clinical Nutrition*, 79, 427-443.
- Burke, L. M., Angus, D. J., Cox, G. R., Cummings, N. K., Febbraio, M. A., Gawthorn, K., Hawley, J. A., Minehan, M., Martin, D. T. and Hargreaves, M. (2000). Effect of fat adaptation and carbohydrate restoration on metabolism and performance during prolonged cycling. *Journal of Applied Physiology*, 89, 2413-2421.
- Burke, L. M., Kiens, B. and Ivy, J. L. (2004). Carbohydrates and fat for training and recovery. *Journal of Sports Sciences*, 22, 15-30.
- Burke, L. M., Wood, C., Pyne, D. B., Telford, R. D. and Saunders, P. U. (2005). Effect of carbohydrate intake on half-marathon performance of well-

trained runners. *International Journal of Sport Nutrition and Exercise Metabolism*, 15, 573-589.

Burke, L. M. and Kiens, B. (2006). "Fat adaptation" for athletic performance - the nail in the coffin? *Journal of Applied Physiology*, 100, 7-8.

Burke, L.M. (2007). Middle- and long-distance running. In *Practical Sports Nutrition*, pp. 109-140. Champaign, Illinois, Human Kinetics.

Bussau, V. A., Fairchild, T. J., Rao, A., Steele, P. D. and Fournier, P. A. (2002). Carbohydrate loading in human muscle: an improved 1 day protocol. *European Journal of Applied Physiology*, 87, 290-295.

Calloway DH. (1975). Nitrogen balance of men with marginal intakes of protein and energy. *Journal of Nutrition* 105, 914-923.

Deakin, V. (2006). Iron depletion in athletes. In: *Clinical Sports Nutrition*, 3<sup>rd</sup> edition (edited by L. Burke & V. Deakin), pp. 263-312, Sydney: McGraw-Hill.

Dennis, S. C. and Noakes, T. D. (1999). Advantages of a smaller bodymass in humans when distance-running in warm, humid conditions. *European Journal of Applied Physiology and Occupational Physiology*, 79, 280-284.

Fogelholm, M. (1995). Inadequate iron status in athletes: an exaggerated problem? In: *Sports nutrition: minerals and electrolytes*. (edited by C. V. Kies & J. A. Driskell), pp. 81-95, Boca Raton: CRC Press.

Forslund AH, El-Khoury AE, Olsson RM, Sjodin AM, Hambraeus L & Young VR. (1999). Effect of protein intake and physical activity on 24-h pattern and rate of macronutrient utilization. *American Journal of Physiology* 276, E964-976.

Friedman JE & Lemon PW. (1989). Effect of chronic endurance exercise on retention of dietary protein. *International Journal of Sports Medicine* 10, 118-123.

Gordon, B., Kohn, L. A., Levine, S. A., Matton, M., Scriver, W. d. and Whiting, W. B. (1925). Sugar content of the blood in runners following a marathon race, with especial reference to the prevention of hypoglycemia: further observations. *Journal of the American Medical Association*, 85, 508-509.

Graham TE. (2001). Caffeine and exercise: metabolism, endurance and performance. *Sports Med* 31, 785-807.

Hargreaves, M. (1999). Metabolic responses to carbohydrate ingestion: effects on exercise performance. In: *Perspectives in exercise science and sports medicine*, Vol. 12: The metabolic basis of performance in exercise and sport (edited by D. R. Lamb & R. Murray), pp. 93-124). Carmel, In.: Cooper.

Havemann, L., West, S., Goedecke, J.H., McDonald, I.A., St-Clair Gibson, A., Noakes, T.D., and Lambert, E.V. (2006). Fat adaptation followed by carbohydrate-loading compromises high-intensity sprint performance. *Journal of Applied Physiology*, 100, 194-202.

Hawley, J.A., Gibala, M.J., and Berman, S. (2007). Innovations in athletic preparation: role of substrate availability to modify training adaptation and performance. *Journal of Sports Sciences*

- Karlsson, J. and Saltin, B. (1971). Diet, muscle glycogen, and endurance performance. *Journal of Applied Physiology*, 31, 203-206.
- Klingshirn, L. A., Pate, R. R., Bourque, S. P., Davis, J. M. and Sargent, R. G. (1992). Effect of iron supplementation on endurance capacity in iron-depleted female runners. *Medicine and Science in Sports and Exercise*, 24, 819-824.
- Kirwan, J. P., Costill, D. L., Mitchell, J. B., Houmard, J. A., Flynn, M. G., Fink, W. J. and Beltz, J. D. (1988). Carbohydrate balance in competitive runners during successive days of intense training. *Journal of Applied Physiology*, 65, 2601-2606.
- Lamanca, J. J. and Haymes, E. M. (1993). Effects of iron repletion on VO<sub>2</sub>max, endurance, and blood lactate in women. *Medicine and Science in Sports and Exercise*, 25, 1386-1392.
- Levine, S. A., Gordon, B. and Derick, C. L. (1924). Some changes in the chemical constituents of the blood following a marathon race, with special reference to the development of hypoglycemia. *Journal of the American Medical Association*, 82, 1778-1779.
- Lucia, A., Esteve-Lanao, J., Olivan, J., Gomez-Gallego, F., San Juan, A. F., Santiago, C., Perez, M., Chamorro-Vina, C., and Foster, C. (2006). Physiological characteristics of the best Eritrean runners-exceptional running economy. *Applied Physiology: Nutrition and Metabolism*, 31, 530-540.
- McKenzie, S., Phillips, S., Carter, S.L., Lowther, S., Gibala, M.J. and Tarnopolsky, M.A. (2000). Endurance exercise training attenuates leucine oxidation and branched-chain 2-oxo acid dehydrogenase activation during exercise in humans. *American Journal of Physiology*, 278, E580–E587.
- Manore, M.M., Kam, L.C., Loucks, A.B. (2007) The female athlete triad: components, nutrition issues, and health consequences. *Journal of Sports Sciences*
- Marino, F. E., Mbambo, Z., Kortekaas, E., Wilson, G., Lambert, M. I., Noakes, T. D. and Dennis, S. C. (2000). Advantages of smaller body mass during distance running in warm, humid environments. *Pflügers Archives - European Journal of Physiology*, 441, 359-367.
- Maughan, R.J., Depiesse, F. and Geyer, H. (2007) The use of dietary supplements by athletes. *Journal of Sports Sciences*
- Meredith CN, Zackin MJ, Frontera WR & Evans WJ. (1989). Dietary protein requirements and body protein metabolism in endurance-trained men. *Journal of Applied Physiology*, 66, 2850-2856.
- Meyer, F., O`Connor, H., Shirreffs, S.M. (2007). Nutrition for the young athlete. *Journal of Sports Sciences*.
- Millard-Stafford, M. L., Sparling, P. B., Roszkopf, L. B. and Dicarlo, L. J. (1992). Carbohydrate-electrolyte replacement improves distance running performance in the heat. *Medicine and Science in Sports and Exercise*, 24, 934-940.
- Millard-Stafford, M., Roszkopf, L. B., Snow, T. K. and Hinson, B. T. (1997). Water versus carbohydrate-electrolyte ingestion before and during a 15-km run in the heat. *International Journal of Sport Nutrition*, 7, 26-38.

Nelson, A. G., Arnall, D. A., Kokkonen, J., Day, R. and Evans, J. (2001). Muscle glycogen supercompensation is enhanced by prior creatine supplementation. *Medicine and Science in Sports and Exercise*, 33, 1096-1100.

Nielsen, P. and Nachtigall, D. (1998). Iron supplementation in athletes: current recommendations. *Sports Medicine*, 26, 207-216.

Nimmo, M.A., and Ekblom, B. (2007). Fatigue and illness in athletes. *Journal of Sports Sciences*

Noakes, T. D. (2002). IMMDA advisory statement of guidelines for fluid replacement during marathon running. *New studies in athletics: the IAAF technical quarterly*, 17, 15-24.

O'Connor, H., Olds, T., and Maughan, R.J. 2007 Physique & performance for track & field events. *Journal of Sports Sciences*

Oopik, V., Saaremets, I., Medijainen, L., Karelson, K., Janson, T. and Timpmann, S. (2003). Effects of sodium citrate ingestion before exercise on endurance performance in well-trained runners. *British Journal of Sports Medicine*, 37, 485-489.

Oopik, V., Saaremets, I., Timpmann, S., Medijainen, L. and Karelson, K. (2004). Effects of acute ingestion of sodium citrate on metabolism and 5 km running performance: a field study. *Canadian Journal of Applied Physiology*, 29, 691-703.

Phillips, S.M., Atkinson, S.A., Tarnopolsky, M.A., and MacDougall, J.D (1993). Gender differences in leucine kinetics and nitrogen balance in endurance athletes. *Journal of Applied Physiology*, 75, 2134-2141

Pitsiladis, Y. P., Duignan, C. and Maughan, R. J. (1996). Effects of alterations in dietary carbohydrate intake on running performance during a 10 km treadmill time trial. *British Journal of Sports Medicine*, 30, 226-231.

Reilly, T., Waterhouse, J., Burke, L.M., and Alonso, J.M. (2007) Nutrition for travel. *Journal of Sports Sciences*

Riddell MC, Partington SL, Stupka N, Armstrong D, Rennie C & Tarnopolsky MA.

(2003). Substrate utilization during exercise performed with and without glucose ingestion in female and male endurance trained athletes. *International Journal of Sport Nutrition and Exercise Metabolism* 13, 407-421.

Roy BD, Luttmer K, Bosman MJ & Tarnopolsky MA. (2002). The influence of post-exercise macronutrient intake on energy balance and protein metabolism in active females participating in endurance training. *International Journal of Sport Nutrition and Exercise Metabolism* 12, 172-188.

Saltin, B., Larsen, H., Terrados, N., Bangsbo, J., Bak, T., Kim, C. K., Svedenag, J., and Rolf, C. J. (1995). Aerobic exercise capacity at sea level and at altitude in Kenyan boys, junior and senior runners compared with Scandinavian runners. *Scandinavian Journal of Medicine and Science in Sports*, 5, 209-221.

Schoene, R. B., Escourrou, P., Robertson, H. T., Nilson, K. L., Parsons, J. R. and Smith, N. J. (1983). Iron repletion decreases maximal exercise lactate concentrations in female athletes with minimal iron-deficiency anemia. *Journal of Laboratory and Clinical Medicine*, 102, 306-312.

Seiler, K. S., and Kjerland, G. O. (2006). Quantifying training intensity distribution in elite endurance athletes: is there evidence for an "optimal" distribution? *Scandinavian Journal of Medicine and Science in Sports*, 16, 49-56.

Sherman, W. M., Costill, D. L., Fink, W. J. and Miller, J. M. (1981). Effect of exercise-diet manipulation on muscle glycogen and its subsequent utilisation during performance. *International Journal of Sports Medicine*, 2, 114-118.

Sherman, W. M., Costill, D. L., Fink, W. J., Hagerman, F. C., Armstrong, L. E. and Murray, T. F. (1983). Effect of a 42.2 km footrace and subsequent rest or exercise on muscle glycogen and enzymes. *Journal of Applied Physiology*, 55, 1219-1224.

Sherman, W. M., Doyle, J. A., Lamb, D. R. and Strauss, R. H. (1993). Dietary carbohydrate, muscle glycogen, and exercise performance during 7 d of training. *American Journal of Clinical Nutrition*, 57, 27-31.

Shirreffs, S.M., Casa, D.J., Carter, R. 2007. Fluid needs for training and competition in athletics. *Journal of Sports Sciences*

Sullo, A., Monda, A., Brizzi, G., Meninno, V., Papa, A., Lombardo, P. and Fabbri, B. (1998). The effect of a carbohydrate loading on running performance during a 25-km treadmill time trial by level of aerobic capacity. *European Review for Medical and Pharmacological Sciences*, 2, 195-202.

Tarnopolsky MA. (2000). Gender differences in substrate metabolism during endurance exercise. *Canadian Journal of Applied Physiology* 25, 312-327.

Tarnopolsky MA, Atkinson SA, Phillips SM & MacDougall JD. (1995). Carbohydrate loading and metabolism during exercise in men and women. *Journal of Applied Physiology* 78, 1360-1368.

Tarnopolsky MA, Bosman M, Macdonald JR, Vandeputte D, Martin J & Roy BD. (1997). Postexercise protein-carbohydrate and carbohydrate supplements increase muscle glycogen in men and women. *Journal of Applied Physiology* 83, 1877-1883.

Tarnopolsky MA, MacDougall JD & Atkinson SA. (1988). Influence of protein intake and training status on nitrogen balance and lean body mass. *Journal of Applied Physiology* 64, 187-193.

Tarnopolsky MA, Zawada C, Richmond LB, Carter S, Shearer J, Graham T & Phillips SM. (2001). Gender differences in carbohydrate loading are related to energy intake. *Journal of Applied Physiology* 91, 225-230.

Tsintzas, K., Liu, R., Williams, C., Campbell, I. and Gaitanos, G. (1993). The effect of carbohydrate ingestion on performance during a 30 km race. *International Journal of Sport Nutrition*, 3, 127-139.

Tsintzas, O. K., Williams, C., Singh, R. and Wilson, W. (1995). Influence of carbohydrate-electrolyte drinks on marathon running performance. *European Journal of Applied Physiology and Occupational Physiology*, 70, 154-160.

Tsintzas, O. K., Williams, C., Boobis, L. and Greenhaff, P. (1996a). Carbohydrate ingestion and single muscle fiber glycogen metabolism during prolonged running in men. *Journal of Applied Physiology*, 81, 801-809.

Tsintzas, O. K., Williams, C., Wilson, W. and Burrin, J. (1996b). Influence of carbohydrate supplementation early in exercise on endurance running capacity. *Medicine and Science in Sports and Exercise*, 28, 1373-1379

van Loon, L. J. C., Murphy, R., Oosterlaar, A. M., Cameron-Smith, D., Hargreaves, M., Wagenmakers, A. J. M. and Snow, R. (2004). Creatine supplementation increases glycogen storage but not GLUT-4 expression in human skeletal muscle. *Clinical Science*, 106, 99-106.

Van Nieuwenhoven, M. A., Brouns, F. and Kovacs, E. M. R. (2005). The effect of two sports drinks and water on GI complaints and performance during an 18-km run. *International Journal of Sports Medicine*, 26, 281-285.

Wallis GA, Dawson R, Achten J, Webber J & Jeukendrup AE. (2006).  
Metabolic  
response to carbohydrate ingestion during exercise in males and females.  
*American Journal of Physiology* 290, E708-715.

Williams, C., Brewer, J. and Walker, M. (1992). The effect of a high carbohydrate diet on running performance during a 30-km treadmill time trial. *European Journal of Applied Physiology and Occupational Physiology*, 65, 18-24.